

You have seen the derivation of the equations for circular motion and solved problems by using them. However, it is always hard to accept a theoretical concept until you test it yourself. In this investigation, you will obtain experimental data for uniform circular motion and compare your data to the theory.

BACKGROUND

In this activity you will be looking at how well the equation $F_c = 4\pi^2 m r f^2$ describes actual experimental results for an object experiencing uniform circular motion.

NOTE!

You will vary 3 different variables:


- tension due to a hanging mass (F_g)
- radius of rotation (r)
- mass of stoppers (m)

INSTRUCTIONS

A. Refer to INV.3.3.2 on page 136 and 137 of your textbook (Nelson Physics 12). Follow procedure steps 1-13.

NOTE!

- ① To obtain reasonable values, you will need to make the measurements several times and take the average.
- ② Record the time and frequency to 2 decimal places and the remaining values to 3 significant digits.
- ③ Use the data table on the next page to record your data. Recall: $f = N/t$
- ④ Use the equation $4\pi^2 m r f^2$ to calculate an experimental value of F_c for each radii.
- ⑤ Calculate the percentage error between the theoretical (accepted) and experimental (measured) centripetal forces.



QUESTIONS

1. What sources of error did you encounter? How did they affect your results? How did you minimize them? To improve your results, what would you do differently next time?
2. To obtain the best accuracy, the tension force acting on the stopper should be horizontal. What should happen to the accuracy as the frequency of revolution of the stopper(s) increases? Explain your reasoning.
3. Use the equation $4\pi^2 m r f^2$ to answer the following questions.
 - (a) Determine the relationship between F_c and (i) m , (ii) r , and (iii) f . Express your answer as a proportionality.
 - (b) What happens to F_c when (i) m , (ii) r , and (iii) f are halved? Express your answer as a multiplier.
 - (c) How can you keep F_c and r constant if m doubles? if f triples? Explain your reasoning with the aid of a proportionality statement.
4. A conical pendulum swings in a circle, as shown. Show that the form of the equation relating the angle that the string makes with the vertical to the speed of the bob is identical to the equation for the banking of curves. (i.e. $\tan\theta = v^2/rg$)

- L = length of pendulum
 θ = angle with vertical
 F_T = tension in the string
 mg = weight of bob (i.e. F_g)
 r = radius of circular path
 h = height of object from ceiling

